



NCERT



CHAPTER WISE TOPIC WISE

LINE BY LINE QUESTIONS

2024



BY
SCHOOL OF
EDUCATORS

- Based on Physical State**
- Gas in Gas (Air)
 - Gas in liquid (Soda water)
 - Gas in Solid (Hydrogen in Pd)
 - Liquid in gas (fog)
 - Liquid in liquid (alcohol in H_2O)
 - Liquid in Solid (Amalgams)
 - Solid in gas (smog)
 - Solid in liquid (sugar in water)
 - Solid in Solid (alloys)



- Based on Concentration**
- Dilute Solution
 - Concentrated Solution
 - Saturated Solution
 - Super Saturated Solution



Types of Solutions

SOLUBILITY

Gas in liquid

Effect of pressure

Solubility increases with increase in pressure

Henry's Law

Partial pressure of gas in vapour phase
 $P = K_H X$

Solid in liquid

Effect of nature of solute of solvent
→ Like dissolves like


Effect of temp →
Exothermic process → Increase with rise in temp.
Effect → Endothermic process → Decrease with rise in temp.
Effect of pressure → No effect

Effect of pressure

Raoult's Law

Partial vapour pressure = Partial pressure × Mole fraction
 $P_j = P_j^0 X_j + P_j^0 X_B$

Homogeneous mixture of two or more pure substance



Concentration terms



Mole fraction (x)

$$= \frac{\text{No. of moles of solute}}{\text{Total moles in solution}}$$

Mass percentage (w/w)

$$= \frac{\text{Mass of solute}}{\text{Total mass of solution}} \times 100$$

Parts per million (PPM)

$$= \frac{\text{No. of parts of solute}}{\text{Total no. of parts of all components of solution}}$$

Volume % (V/V)

$$= \frac{\text{Volume of solute}}{\text{Total volume of solution}} \times 100$$

$$\text{Mass by volume \%} = \frac{\text{mass of solute}}{\text{Total mass in solution}} \times 100$$

Normality (N)

$$= \frac{\text{No. of grams equivalent of solute}}{\text{Volume of solution (L)}}$$

Molality (M)

$$= \frac{\text{No. of moles of solute}}{\text{Weight of the solvent in kg}}$$

Molarity (M)

$$= \frac{\text{No. of moles of solute}}{\text{Volume of solution (L)}}$$

Colligative Property

Osmotic pressure
Pressure applied to stop the flow of solvent through semi permeable membrane
 $\pi = CRT$

$$\Delta T_b = \frac{K_b \times 1000 \times w_2}{M_2 \times w_1}$$

$$\Delta T_f = \frac{K_f \times w_2 \times 1000}{M_2 \times w_1}$$

$$\frac{w_2 \times M_1}{M_2 \times w_1} = \frac{P_2 - P_s}{P_s}$$

Van't Hoff factor (i)


Ratio of the normal mass to the observed molecular mass of the solute:
 $i = \frac{\text{Normal molar mass}}{\text{Observed molar mass}}$

Ideal & Non Ideal Solution


Does not obey Raoult's law
 $P_j \neq P_j^0 X_j + P_j^0 X_B$

Ideal


$$P_j = P_j^0 X_j + P_j^0 X_B$$



Positive deviation



Negative deviation



Azeotropes

Constant boiling mixtures

Positive deviation mixture
Minimum boiling azeotropes
Ex. 95% Ethanol in water

Negative deviation mixture
Maximum boiling azeotropes
Ex. 68% HNO_3 in water

NCERT LINE BY LINE QUESTIONS


- (1.)** Which one of the following electrolytes has the same value of Van't Hoff factor (i) as that of $\text{Al}_2(\text{SO}_4)_3$ (if all are 100% ionized)
- (a.) $\text{K}_4[\text{Fe}(\text{CN})_6]$ (b.) K_2SO_4
 (c.) $\text{K}_3[\text{Fe}(\text{CN})_6]$ (d.) $\text{Al}(\text{NO}_3)_3$
- (2.)** Maximum amount of a solid solute that can be dissolved in a specific amount of a given liquid solvent does not depend upon—
- (a.) temperature (b.) nature of solute
 (c.) pressure (d.) nature of solvent
- (3.)** If solute and solvent interactions are more than solute-solute and solvent-solvent interaction then
- (a.) it is ideal solution. (b.) it is non-ideal solution with position deviation.
 (c.) it is non-ideal solution with negative deviation. (d.) can't be predicted.
- (4.)** When 40 g of substance is dissolved in 1000 g of water, its freezing point is depressed by 1.86°C . K_f for water is $1.86^\circ\text{C mol}^{-1}$ then find the molar mass of the solute
- (a.) 4 (b.) 10
 (c.) 40 (d.) 400
- (5.) Assertion:** When a solution is separated from the pure solvent by a semipermeable membrane, the solvent molecules pass through it from pure solvent side to the solution side.
Reason: Diffusion of solvent occurs from a region of high concentration solution to a region of low concentration solution.
- (a.) Both A and R are true and R is the correct explanation of A. (b.) Both A and R are true but R is not the correct explanation of A.
 (c.) A is true but R is false. (d.) Both A and R are false.
- (6.)** In water saturated air the mole fraction of water vapour is 0.02. If the total pressure of the saturated air is 1.2 atm, the partial pressure of dry air is
- (a.) 1.18 atm (b.) 1.76 atm
 (c.) 1.176 atm (d.) 0.98 atm
- (7.)** Matrix match.

Column 1	Column 2
(A) Standard	(p) Solution with two solution components
(B) Solid solution	(q) A solution which contains maximum amount of solute that can be dissolved in a given amount of solvent at given temperature
(C) Binary solution	(r) A solution in solid phase

- (a.) $A \rightarrow P, B \rightarrow Q, C \rightarrow R$ (b.) $A \rightarrow Q, B \rightarrow R, C \rightarrow P$



- (c.) $A \rightarrow R, B \rightarrow P, C \rightarrow Q$ (d.) $A \rightarrow R, B \rightarrow Q, C \rightarrow P$
- (8.) According to Henry law, the amount of gas that will dissolve in blood plasma or any other liquid is determined by which of these factors
- (a.) Solubility of the gas in the liquid (b.) The total pressure of the gas mixture
- (c.) pH of the liquid (d.) The osmotic pressure of the gas mixture
- (9.) The freezing point depression constant for water is $-1.86^\circ\text{C}\cdot\text{mol}^{-1}$. If 5.0 g Na_2SO_4 is dissolved in 45 g H_2O , the freezing point is changed by -3.82°C . Calculate Van't Hoff factor for Na_2SO_4 :
- (a.) 2.05 (b.) 2.63
- (c.) 3.11 (d.) 0.381
- (10.) vapour phase diagram for a solution is given below if dotted lines represent deviation. The correct observation for this solution
- (a.) ΔH_{mix} : positive (b.) ΔS_{mix} : positive
- (c.) ΔV_{mix} : positive (d.) All of these
- (11.) Osmotic pressure of a solution at a given temperature
- (a.) increase with concentration (b.) decrease with concentration
- (c.) remain same (d.) initially increase and then decrease
- (12.) Read statements
- I. Semipermeable membrane contains network of submacroscopic holes or pores.
- II. Semipermeable membrane appears to be continuous sheets or films.
- III. Solvent molecule cannot pass through the holes of semipermeable membrane but solute molecule can pass.
- On the basis of the statements given above select correct option
- (a.) Only (I) (b.) Only (II)
- (c.) Only (III) (d.) All of these
- (13.) The vapour of a solution having 2.0 g of solute X (molar atomic mass = 32 g mol^{-1}) in 100 g of CS_2 (vapour pressure = 854 torr) is 848.9 torr . The molecular formula of solute is
- (a.) X (b.) X_2
- (c.) X_4 (d.) X_8
- (14.) The Van't Hoff factor of 0.1M $\text{Ba}(\text{NO}_3)_2$ solution is 2.74. The degree of dissociation is
- (a.) 91.3% (b.) 87%
- (c.) 100% (d.) 74%
- (15.) The vapour pressure of two liquids P and Q are 80 and 60 torr respectively. The total vapour pressure of solution obtained by mixing 3 mole of P and 2 mole of Q would be
- (a.) 72 torr (b.) 140 torr
- (c.) 68 torr (d.) 20 torr
- (16.) The units of ebullioscopic constant is

- (a.) $\text{K kg}^{-1}\text{mol}^{-1}$ or $\text{K}(\text{molality})$ (b.) mol kg K^{-1} or $\text{K}^{-1}(\text{molality})$
 (c.) $\text{kg mol}^{-1}\text{K}^{-1}$ or $\text{K}^{-1}(\text{molality})$ (d.) K mol kg^{-1} or $\text{K}(\text{molality})$
- (17.)** Of the following 0.10 m aqueous solutions, which one will exhibit the largest freezing point depression?
 (a.) KCl (b.) $\text{C}_6\text{H}_{12}\text{O}_6$
 (c.) $\text{Al}_2(\text{SO}_4)_3$ (d.) K_2SO_4
- (18.)** An aqueous solution of sugar is taken in a beaker. At freezing point of solution ,
 (a.) Crystals of sugar separated (b.) Crystals of glucose and fructose are separated
 (c.) Crystals of sucrose separated (d.) Mixture of ice and some sugar crystals separated
- (19.)** Which one of the following binary mixtures forms an azeotrope with minimum boiling type
 (a.) acetone— ethanol (b.) $\text{H}_2\text{O} - \text{HNO}_3$
 (c.) benzene— toluene (d.) n-hexane—n-heptane
- (20.)** How many grams of CH_3OH should be added to water to prepare 150 ml solution of 2M CH_3OH
 (a.) 9.6×10^3 (b.) 2.4×10^3
 (c.) 9.6 (d.) 2.4
- (21.)** Relationship between partial pressure & mole fraction is
 (a.) Graham's law (b.) Raoult's law
 (c.) Le-Chatelier (d.) Avogadro's law
- (22.)** 4L of 0.02M aqueous solution NaCl was diluted by adding one litre of water. The molality of resultant solution is—
 (a.) 0.004 (b.) 0.008
 (c.) 0.012 (d.) 0.016
- (23.)** Iodine and sulphur dissolve in
 (a.) H_2O (b.) benzene
 (c.) CS_2 (d.) ethanol
- (24.)** 19.5 g of CH_2FCOOH is dissolved in 500 g of water. The depression in the freezing point of water observed is 1.0°C . Calculate Van't Hoff factor.
 (a.) 2.97 (b.) 0.79
 (c.) 6.28 (d.) 1.0753
- (25.)** Consider the figure



Mark the correct statement

- (a.) water will move from side (A) to side (B) if a pressure lower than osmotic pressure is applied on piston (B).
 (b.) Water will move from side (B) to side (A) if a pressure greater than osmotic pressure is applied on piston (B).
 (c.) The value of molal depression constant depends on nature of solvent.
 (d.) Relative lowering of vapour pressure is a dimensionless quantity.
- (26.) P_A and P_B are the vapour pressure of pure liquid components A and B respectively of an ideal binary solution. If X_A represent the mole fraction of component A, the total pressure of the solution will be —
 (a.) $P_A + X_A (P_B - P_A)$
 (b.) $P_A + X_A (P_A - P_B)$
 (c.) $P_B + X_A (P_B - P_A)$
 (d.) $P_B + X_A (P_A - P_B)$
- (27.) Nalorphene ($C_{19}H_{21}NO_3$), similar to morphine, is used to combat withdrawal symptoms in narcotic users. Dose of nalorphene generally given is 1.5 mg. Calculate the mass of 1.5×10^{-3} m aqueous solution required for the above dose
 (a.) 9.2 g
 (b.) 4.2 g
 (c.) 3.2 g
 (d.) 6.2 g
- (28.) **Assertion:** Azeotropic mixture are formed only by non-ideal solution.
Reason: Boiling point of an azeotropic mixture is either higher or lower than both the components of solution.
 (a.) Both A and R are correct and R is correct explanation of A.
 (b.) Both A and R are correct but R is not correct explanation of A.
 (c.) A is true but R is false.
 (d.) Both A and R are false.
- (29.) If molality of a dilute solution is double, the value of molal depression constant (K_f) will be?
 (a.) Halved
 (b.) tripled
 (c.) unchanged
 (d.) doubled
- (30.) Solid solution in which the solute is gas
 (a.) copper dissolved in gold
 (b.) camphor in nitrogen gas
 (c.) solution of hydrogen in palladium
 (d.) all of the above
- (31.) Which of the following 0.10m aqueous solutions will have the lowest freezing point?



(32.) Which one is a colligative property?

(a.) Boiling point

(b.) Vapour pressure

(c.) Osmotic pressure

(d.) Freezing point

(33.) H_2S , a toxic gas with rotten egg like smell, is used for the qualitative analysis. If the solubility of H_2S in water at STP is 0.195m. Calculate Henry law constant?

(a.) 282 bar

(b.) 386 bar

(c.) 192 bar

(d.) 465 bar

(34.) If liquids A and B form an ideal solution.

(a.) the free energy of mixing is zero

(b.) the free energy as well as entropy of mixing are zero

(c.) enthalpy of mixing is zero

(d.) the entropy of mixing is zero

(35.) 200cm^3 of an aqueous solution of a protein contains 1.26 g of the protein. The osmotic pressure of such solution at 300 K is found to be $2.57 \times 10^{-3}\text{bar}$. Calculate the molar mass of protein.

(a.) 21200 g mol^{-1}

(b.) 61022 g mol^{-1}

(c.) 19200 g mol^{-1}

(d.) none of these

(36.) Low concentration of oxygen in the blood and tissues of people living at high altitude is due to

(a.) Low temperature

(b.) Low atmospheric pressure

(c.) High atmospheric pressure

(d.) Both low temperature and high atmospheric pressure

(37.) Which of the following statement(s) is/are true

I. In a binary mixture, mole fraction of A is

$$X_A = \frac{n_A}{n_A + n_B}$$

II. For solution containing i number of components, $X_i = \frac{n_i}{\sum n_i}$

III. Sum of all mole fraction is one.

Select option with true statements.

(a.) I and II

(b.) II and III

(c.) III and I

(d.) All of the above

(38.) When a solute is present in trace quantities which of the following expression is used?

(a.) Gram per million

(b.) Milligram percent

(c.) Microgram percent

(d.) Parts per millions

(39.) Which of the following physical property is used to determine the molecular mass of a polymer Solution?

(a.) Relative lowering of vapour pressure

(b.) Elevation in boiling point

- (c.) Depression in freezing point (d.) Osmotic pressure

(40.) The freezing point of 1% solution of lead nitrate in water will be

- (a.) 2°C (b.) 1°C
(c.) 0°C (d.) below 0°C

(41.) If 0.15 g of solute, dissolved in 15 g of solvent, is boiled at a temperature higher by 0.216°C than that of pure solvent, the molecular weight of substance (molal elevation constant for the solvent is 2.16°C) is

- (a.) 10.1 (b.) 100
(c.) 1.01 (d.) 1000

(42.) Which condition is not satisfied by an ideal solution .

- (a.) $\Delta_{\text{mix}} V = 0$ (b.) $\Delta_{\text{mix}} S = 0$
(c.) obedience to Raoult's law (d.) $\Delta_{\text{mix}} H = 0$


(43.) How many grams of Conc. HNO_3 solution should be used to prepare 250 ml of $2.0\text{M}\text{HNO}_3$? The conc.acid is 70 % HNO_3 ?

- (a.) 70 g conc. HNO_3 (b.) 54 g conc. HNO_3
(c.) 45 g conc. HNO_3 (d.) 90 g conc. HNO_3

(44.) 0.5 molal aqueous solution of a weak acid (HX) is 20% ionised. If κ_f for water is $1.86\text{ K kg mol}^{-1}$. The lowering freezing point of the solution is

- (a.) 0.56 K (b.) 1.12 K
(c.) -0.56K (d.) -1.12K

(45.) The freezing point of solution is marked as:



- (a.) A (b.) B
(c.) C (d.) D

(46.) Which condition is not satisfied by an ideal solution?

- (a.) $\Delta_{\text{mix}} H = 0$ (b.) $\Delta_{\text{mix}} V = 0$

- (c.) $\Delta_{\text{mix}} S = 0$ (d.) obedience to Raoult's law
- (47.) The relative lowering of the vapour pressure is equal to the ratio between number of
- (a.) solute molecules to the solvent molecules (b.) solute molecules to the total molecules in the solution
- (c.) solvent molecules to the total molecules in the solution (d.) solvent molecules to the total number of ions of the solute
- (48.) If solubility of any gas in the liquid at 1 bar pressure is 0.05 mol / litre, what will be its solubility at 3 bar pressure, keeping the temperature constant?
- (a.) $\frac{0.05}{3}$ mol / L (b.) 0.15 mol / L
- (c.) 0.05 mol / L (d.) 1.0 mol / L
- (49.) **Assertion:** Camphor is used as a solvent in the experimental determination of molecular masses of naphthalene and anthracene.
Reason: Camphor has high molal elevation constant.
- (a.) Both A and R are correct and R is the correct explanation of A. (b.) Both A and R are correct and R is not correct explanation of A.
- (c.) A is correct but R is incorrect statement. (d.) Both A and R are false.
- (50.) At equilibrium, the rate of dissolution of a solid solute in a volatile liquid solvent is —
- (a.) less than rate of crystallization (b.) greater than rate of crystallization
- (c.) equal to the rate of crystallization (d.) zero

TOPIC WISE PRACTICE QUESTIONS

TOPIC 1: Solubility and Concentration of Solutions

- When the solute is present in trace quantities the following expression is used
 1) gram per million 2) milligram percent
 3) microgram percent 4) parts per million
- Which one of the following gases has the lowest value of Henry's law constant?
 1) N_2 2) He 3) H_2 4) CO_2
- Equal moles of water and urea are taken in a flask. What is mass percentage of urea in the solution ?
 1) 7.692% 2) 9.2% 3) 76.92% 4) 0.7692%
- What is the normality of a 1 M solution of H_3PO_4 ?
 1) 0.5 N 2) 1.0 N 3) 2.0 N 4) 3.0 N
- Molarity of liquid HCl will be, if density of solution is 1.17 gm/cc
 1) 36.5 M 2) 32.05 M 3) 18.25 M 4) 42.10 M
- An X molal solution of a compound in benzene has mole fraction of solute equal to 0.2. The value of X is
 1) 14 2) 3.2 3) 1.4 4) 2
- 1 M, 2.5 litre NaOH solution is mixed with another 0.5 M, 3 litre NaOH solution. Then find out the molarity of resultant solution
 1) 0.80 M 2) 1.0 M 3) 0.73 M 4) 0.50 M

8. In acidic medium, the equivalent weight of $K_2Cr_2O_7$ (Mol. wt. = M) is
 1) M 2) M/2 3) M/3 4) M/6
9. The mole fraction of the solute in one molal aqueous solution is:
 1) 0.009 2) 0.018 3) 0.027 4) 0.036
10. 10 g of NaCl is dissolved in 106g of the solution. Its concentration is
 1) 100 ppm 2) 0.1 ppm 3) 1 ppm 4) 10 ppm
11. Which of the following substances will lose its solubility with increase in temperature?
 1) NaOH 2) Na_2CO_3 3) Na_2SO_4 4) All
12. 2.5 litres of NaCl solution contain 5 moles of the solute. What is the molarity?
 1) 5 molar 2) 2 molar 3) 2.5 molar 4) 12.5 molar
13. Which of the following factor do not affect solubility of solid solute in liquid?
 1) Temperature 2) Pressure 3) Nature of solute 4) All of these

TOPIC 2: Vapour Pressure, Laws of Solutions and Ideal, Non-Ideal Solutions

14. For a dilute solution, Raoult's law states that
 1) the lowering of vapour pressure is equal to the mole fraction of solute.
 2) the relative lowering of vapour pressure is equal to the mole fraction of solute.
 3) the relative lowering of vapour pressure is proportional to the amount of solute in solution.
 4) the vapour pressure of the solution is equal to the mole fraction of solvent.
15. If p° and p_s are vapour pressures of solvent and its solution, respectively, χ_1 and χ_2 are mole fractions of solvent and solute, respectively, then
 1) $p_s = p^\circ / \chi_2$ 2) $p^\circ - p_s = p^\circ \chi_2$ 3) $p_s = p^\circ \chi_2$ 4) $\frac{p^\circ - p_s}{p_s} = \frac{\chi_1}{\chi_1 + \chi_2}$
16. The normal boiling point of water is 373 K. Vapour pressure of water at temperature T is 19 mm Hg. If enthalpy of vaporisation is 40.67 kJ/mol, then temperature T would be
 (Use : $\log 2 = 0.3$, $R : 8.3 \text{ JK}^{-1} \text{ mol}^{-1}$):
 1) 250 K 2) 291.4 K 3) 230 K 4) 290 K
17. For a binary ideal liquid solution, the total vapour pressure of the solution is given as:
 1) $P_{\text{total}} = P_A^\circ + (P_A^\circ - P_B^\circ) x_B$ 2) $P_{\text{total}} = P_B^\circ + (P_A^\circ - P_B^\circ) x_A$
 3) $P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ) x_A$ 4) $P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ) x_B$
18. Moles of Na_2SO_4 to be dissolved in 12 mole water to lower its vapour pressure by 10 mm Hg at a temperature at which vapour pressure of pure water is 50 mm is:
 1) 1.5 mole 2) 2 mole 3) 1 mole 4) 3 mole
19. Equimolar solutions in the same solvent have
 1) different boiling and different freezing points. 2) same boiling and same freezing points.
 3) same freezing point but different boiling points. 4) same boiling point but different freezing points.
20. The solubility of common salt is 36.0 g in 100 g of water at 20 °C. If systems I, II and III contain 40.0, 36.0 and 20.0 g of the salt added to 100.0 g of water in each case, the vapour pressures would be in the order:
 1) $I < II < III$ 2) $I > II > III$ 3) $I = II > III$ 4) $I = II < III$
21. The vapour pressure of two liquids P and Q are 80 and 60 torr, respectively. The total vapour pressure of solution obtained by mixing 3 mole of P and 2 mole of Q would be
 1) 72 torr 2) 140 torr 3) 68 torr 4) 20 torr
22. A mixture of components A and B will show -ve deviation when
 1) $\Delta V_{\text{mix}} > 0$ 2) $\Delta H_{\text{mix}} > 0$
 3) A-B interaction is weaker than A-A and B-B interactions.
 4) A-B interaction is stronger than A-A and B-B interactions.
23. At the state of dynamic equilibrium, for solute + solvent \square solution.
 1) Rate of dissolution = Rate of unsaturation. 2) Rate of dissolution = Rate of unsaturation.
 3) Rate of dissolution = Rate of saturation 4) Rate of crystallization = Rate of saturation.
24. The value of P° for benzene is 640 mm of Hg. The vapour pressure of solution containing 2.5g substance in 39g benzene is 600mm of Hg the molecular mass of X is –

- 1) 65.25 2) 130 3) 40 4) 80
25. An ideal solution is formed when its components
 1) have no volume change on mixing 2) have no enthalpy change on mixing
 3) both (1) and (2) are correct 4) neither (1) nor (2) is correct
26. For which of the following parameters the structural isomers C_2H_5OH and CH_3OCH_3 would be expected to have the same values?(Assume ideal behaviour)
 1) Boiling points 2) Vapour pressure at the same temperature
 3) Heat of vaporization 4) Gaseous densities at the same temperature and pressure
27. Which one of the following is non-ideal solution
 1) Benzene + toluene 2) *n*-hexane + *n*-heptane
 3) Ethyl bromide + ethyl iodide 4) $CCl_4 + CHCl_3$

TOPIC 3: Colligative Properties and Abnormal Molecular Masses

28. When common salt is dissolved in water
 1) the melting point of the solution increases. 2) the boiling point of solution decreases.
 3) both melting point and boiling point decrease. 4) the boiling point of the solution increases.
29. Camphor is often used in molecular mass determination because
 1) it is readily available. 2) it has a very high cryoscopic constant.
 3) it is volatile. 4) it is solvent for organic substances.
30. The normal boiling point of the solution is the temperature at which the vapour pressure of the solution is –
 1) equal to 1 torr 2) equal to 76 mm Hg 3) equal to 2.0 atm 4) equal to 1 atm
31. When solid SnO_2 is added to an aqueous solution of NaOH, the
 1) vapour pressure is lowered. 2) vapour pressure is raised.
 3) osmotic pressure is increased. 4) boiling point is raised.
32. The vapour pressure of a dilute solution of non-volatile solute is P and the VP of a pure solvent is P° . The lowering of the VP is
 1) + ve 2) – ve 3) P/P° 4) P°/P
33. 12 g of a nonvolatile solute dissolved in 108 g of water produces the relative lowering of vapour pressure of 0.1. The molecular mass of the solute is
 1) 80 2) 60 3) 20 4) 40
34. If a thin slice of sugar beet is placed in concentrated solution of NaCl, then
 1) sugar beet will lose water from its cells. 2) sugar beet will absorb water from solution.
 3) sugar beet will neither absorb nor lose water. 4) sugar beet will dissolve in solution.
35. Which salt shows maximum osmotic pressure in its 1 M solution.
 1) $AgNO_3$ 2) Na_2SO_4 3) $(NH_4)_3PO_4$ 4) $MgCl_2$
36. The osmotic pressure of a sugar solution at $24^\circ C$ is 2.5 atm. The concentration of the solution in mole per litre is
 1) 10.25 2) 1.025 3) 1025 4) 0.1025
37. Which has the maximum osmotic pressure at temperature T ?
 1) 100 mL of 1 M urea solution.
 2) 300 mL of 1 M glucose solution.
 3) Mixture of 100 mL of 1 M urea solution and 300 mL of 1 M glucose solution.
 4) All are isotonic.
38. Which one of the following is a colligative property ?
 1) Boiling point 2) Vapour pressure
 3) Osmotic pressure 4) Freezing point
39. Which one of the following aqueous solutions will exhibit highest boiling point?
 1) 0.015 M urea 2) 0.01 M KNO_3 3) 0.10 M Na_2SO_4 4) 0.015 M glucose
40. When a solution containing non-volatile solute freezes, which equilibrium would exist?
 1) solid solvent \rightleftharpoons liquid solvent 2) solid solute \rightleftharpoons liquid solution
 3) solid solute \rightleftharpoons liquid solvent 4) solid solvent \rightleftharpoons liquid solution
41. Which of the following aqueous solution has minimum freezing point ?

- 1) 0.01 m NaCl 2) 0.005 m C₂H₅OH 3) 0.005 m MgI₂ 4) 0.005 m MgSO₄.
42. A solution containing 1.8 g of a compound (empirical formula CH₂O) in 40 g of water is observed to freeze at -0.465 °C. The molecular formula of the compound is (K_f of water = 1.86 kg K mol⁻¹):
1) C₂H₄O₂ 2) C₃H₆ 3) C₄H₈O₄ 4) C₆H₁₂O₆
43. Freezing point of the following equilibrium,
liquid solvent \rightleftharpoons solid solvent is:
1) $\frac{\Delta H - \Delta G}{T\Delta S}$ 2) $\frac{\Delta H}{\Delta S}$ 3) $\frac{\Delta G}{\Delta S}$ 4) $\frac{\Delta S}{\Delta H}$
44. The freezing point of equimolal aqueous solutions will be highest for:
1) C₆H₅NH₃Cl 2) Ca(NO₃)₂ 3) La (NO₃)₂ 4) C₆H₁₂O₆
45. In a 0.5 molal solution of KCl, KCl is 50% dissociated. The freezing point of solution will be (K_f = 1.86 k kg mol⁻¹):
1) 274.674 K 2) 271.60 K 3) 273 K 4) None of these
46. In a 0.2 molal aqueous solution of a weak acid HX the degree of dissociation is 0.25. The freezing point of the solution will be nearest to: (K_f = 1.86 K kg mol⁻¹)
1) -0.26 °C 2) 0.465 °C 3) -0.48 °C 4) -0.465 °C
47. Which one of the following statements is false ?
1) The correct order of osmotic pressure for 0.01 M aqueous solution of each compound is BaCl₂ > KCl > CH₃COOH > sucrose.
2) Isotonic solutions are those solutions which have the same osmotic pressure.
3) Raoult's law states that the vapour pressure of a component over a solution is proportional to its mole fraction in liquid state.
4) Two sucrose solutions of same molality prepared in different solvents will have the same freezing point depression.
48. 0.1 molal aqueous solution of an electrolyte AB₃ is 90% ionised. The boiling point of the solution at 1 atm is : (K_{b(H₂O)} = 0.52 K kg mol⁻¹)
1) 273.19 K 2) 374.92 K 3) 376.4 K 4) 373.19 K
49. In the case of osmosis, solvent molecules move from :
1) higher vapour pressure to lower vapour pressure.
2) higher concentration to lower concentration.
3) lower vapour pressure to higher vapour pressure.
4) higher osmotic pressure to lower osmotic pressure.
50. Which of the following solutions will have maximum osmotic pressure? (Assume 90% dissociation of each salt):
1) Decinormal aluminium sulphate 2) Decinormal barium chloride solution
3) Decinormal sodium sulphate solution
4) Solution of equal volumes of decinormal barium chloride and decinormal sodium sulphate solutions
51. At 25°C, the highest osmotic pressure is exhibited by 0.1 M solution of
1) CaCl₂ 2) KCl 3) glucose 4) urea
52. Osmotic pressure of blood is 7.40 atm, at 27 °C. Number of moles of glucose to be used per litre for an intravenous injection that is to have same osmotic pressure of blood is:
1) 0.3 2) 0.2 3) 0.1 4) 0.4
53. The freezing point of 1% solution of lead nitrate in water will be
1) 2°C 2) 1°C 3) 0°C 4) below 0°C
54. If the elevation in boiling point of a solution of 10 g of solute (mol. wt. = 100) in 100 g of water is Δ T_b, the ebullioscopic constant of water is
1) 10 2) 10 Δ T_b 3) Δ T_b 4) $\frac{\Delta T_b}{10}$
55. The boiling point of a solution of 0.11 g of a substance in 15 g of ether was found to be 0.1 °C higher than that of pure ether. The molecular weight of the substance will be (K_b = 2.16 °K kg mol⁻¹)
1) 148 2) 158 3) 168 4) 178

56. The rise in the boiling point of a solution containing 1.8 g of glucose in 100 g of solvent is 0.1°C . The molal elevation constant of the liquid is
 1) 0.01 K/m 2) 0.1 K/m 3) 1 K/m 4) 10 K/m
57. An aqueous solution freezes at -0.186°C ($K_f = 1.86$, $K_b = 0.512$) what is the elevation in boiling point?
 1) 0.186°C 2) 0.512°C 3) 0.86°C 4) 0.0512°C
58. At temperature 327°C and concentration C , osmotic pressure of a solution is P , the same solution at concentration $C/2$ and at temperature 427°C shows osmotic pressure of 2 atm , value of P will be
 1) $12/7$ 2) $24/7$ 3) $6/5$ 4) $5/6$
59. The relationship between osmotic pressure at 273 K when 10g glucose (P_1), 10g urea (P_2), and 10g sucrose (P_3) are dissolved in 250 ml of water is
 1) $P_1 > P_2 > P_3$ 2) $P_3 > P_1 > P_2$ 3) $P_2 > P_1 > P_3$ 4) $P_2 > P_3 > P_1$
60. Consider the following statements
 I : Osmosis takes place with increase of entropy.
 II : Osmosis is a non-spontaneous process.
 III : Free energy decreases during osmosis.
 Which of the above is/are correct?
 1) I only 2) I and II 3) II and III 4) I and III
61. For an electrolyte, elevation of B.P. is directly proportional to
 1) molarity 2) molality 3) mole fraction 4) All of these
62. The molecular weight of benzoic acid in benzene as determined by depression in freezing point method corresponds to
 1) ionization of benzoic acid. 2) dimerization of benzoic acid.
 3) trimerization of benzoic acid. 4) solvation of benzoic acid.
63. Blood cells retain their normal shape in solution which are
 1) hypotonic to blood 2) isotonic to blood 3) hypertonic to blood 4) equinormal to blood.
64. Which of the following pairs of solution are isotonic at the same temperature ?
 1) $0.1\text{ M Ca(NO}_3)_2$ and $0.1\text{ M Na}_2\text{SO}_4$ 2) 0.1 M NaCl and $0.1\text{ M Na}_2\text{SO}_4$
 3) 0.1 M urea and 0.1 M MgCl_2 4) 0.2 M urea and 0.1 M NaCl
65. The van't Hoff factor i for a compound which undergoes dissociation in one solvent and association in other solvent is respectively
 1) less than one and greater than one. 2) less than one and less than one.
 3) greater than one and less than one. 4) greater than one and greater than one.
66. Van't Hoff factor is given by the expression _____.
 1) $i = \frac{\text{Normal molar mass}}{\text{Abnormal molar mass}}$ 2) $i = \frac{\text{Abnormal molar mass}}{\text{Normal molar mass}}$
 3) $i = \frac{\text{Observed Colligative property}}{\text{Calculated colligative property}}$ 4) both 1) and 3)
67. If α is the degree of dissociation of Na_2SO_4 , the Vant Hoff's factor (i) used for calculating the molecular mass is
 1) $1 + \alpha$ 2) $1 - \alpha$ 3) $1 + 2\alpha$ 4) $1 - 2\alpha$
68. The correct order of osmotic pressure of 0.01M aqueous solution of the following is
 1) Sucrose $>$ CH_3COOH $>$ KCl 2) CH_3COOH $>$ Sucrose $>$ KCl
 3) Sucrose $>$ KCl $>$ CH_3COOH 4) KCl $>$ CH_3COOH $>$ Sucrose
69. Which one of the following salts will have the same value of van't Hoff factor (i) as that of $\text{K}_4[\text{Fe(CN)}_6]$.
 1) $\text{Al}_2(\text{SO}_4)_3$ 2) NaCl 3) $\text{Al}(\text{NO}_3)_3$ 4) Na_2SO_4 .
70. Consider the following statements
 1. Isotonic solutions have the same molar concentration at a given temperature
 2. The molal elevation constant K_b is a characteristic of a solvent, and is independent of the solute
 added
 3. The freezing point of a 0.1 M aqueous KCl solution is more than that of a 0.1 M aqueous AlCl_3 solution.

Which of these statements is correct?

1) 1 and 2

2) 2 and 3

3) 1 and 3

4) 1, 2 and 3

NEET PREVIOUS YEARS QUESTIONS

- If molality of the dilute solutions is doubled, the value of molal depression constant (K_f) will be : [2017]
1) halved 2) tripled 3) unchanged 4) doubled
- Which of the following statement about the composition of the vapour over an ideal 1 : 1 molar mixture of benzene and toluene is correct? Assume that the temperature is constant at 25°C. (given : Vapour Pressure Data at 25°C, benzene = 12.8 kPa, toluene = 3.85 kPa) [2016]
1) The vapour will contain a higher percentage of benzene
2) The vapour will contain a higher percentage of toluene
3) The vapour will contain equal amounts of benzene and toluene
4) Not enough information is given to make a predication
- At 100 °C the vapour pressure of a solution of 6.5g of a solute in 100 g water is 732 mm. If $K_b = 0.52$, the boiling point of this solution will be [2016]
1) 101 °C 2) 100 °C 3) 102 °C 4) 103 °C
- Which one of the following electrolytes has the same value of van't Hoff's factor (i) as that of the $Al_2(SO_4)_3$ (if all are 100% ionised)? [2015]
1) $K_3[Fe(CN)_6]$ 2) $Al(NO_3)_3$ 3) $K_4[Fe(CN)_6]$ 4) K_2SO_4
- The boiling point of 0.2 mol kg^{-1} solution of X in water is greater than equimolal solution of Y in water. Which one of the following statements is true in this case? [2015]
1) Molecular mass of X is greater than the molecular mass of Y.
2) Molecular mass of X is less than the molecular mass of Y.
3) Y is undergoing dissociation in water while X undergoes no change.
4) X is undergoing dissociation in water.
- Which one is not equal to zero for an ideal solution: [2015]
1) ΔS_{mix} 2) ΔV_{mix} 3) $\Delta P = P_{observed} - P_{Raoult}$ 4) ΔH_{mix}
- Of the following 0.10 m aqueous solutions, which one will exhibit the largest freezing point depression? [2014]
1) KCl 2) $C_6H_{12}O_6$ 3) $Al_2(SO_4)_3$ 4) K_2SO_4
- For an ideal solution, the correct option is :- [2019]
(1) $\Delta_{mix} S = 0$ at constant T and P (2) $\Delta_{mix} V \neq 0$ at constant T and P
(3) $\Delta_{mix} H = 0$ at constant T and P (4) $\Delta_{mix} G = 0$ at constant T and P
- The mixture that forms maximum boiling azeotrope is : [2019]
(1) Water + Nitric acid (2) Ethanol + Water (3) Acetone + Carbon disulphide
(4) Heptane + Octane
- Which of the following statements is correct regarding a solution of two compounds A and B exhibiting positive deviation from ideal behaviour? [2019 ODISSA]
(1) Intermolecular attractive forces between A-A and B-B are stronger than those between A-B.
(2) $\Delta_{mix} H = 0$ at constant T and P (3) $\Delta_{mix} V = 0$ at constant T and P
(4) Intermolecular attractive forces between A-A and B-B are equal to those between A-B.
- In water saturated air the mole fraction of water vapour is 0.02. If the total pressure of the saturated air is 1.2 atm, the partial pressure of dry air is : ? [2019 ODISSA]
(1) 1.18 atm (2) 1.76 atm (3) 1.176 atm (4) 0.98 atm
- The density of 2 M aqueous solution of NaOH is 1.28 g/ cm^3 . The molality of the solution is [Given that molecular mass of NaOH = 40 g mol^{-1}] [2019 ODISSA]

- (1) 1.20 m (2) 1.56 m (3) 1.67 m (4) 1.32 m
13. If 8g of a non-electrolyte solute is dissolved in 114 g of n-octane to reduce its vapour pressure to 80%, the molar mass (in g mol^{-1}) of the solute is [Given that molar mass of n-octane is 114 g mol^{-1}]. [2020 COVID-19]
- (1) 40 (2) 60 (3) 80 (4) 20
14. Isotonic solutions have same [2020 COVID-19]
- (1) vapour pressure (2) freezing temperature (3) osmotic pressure (4) boiling temperature
15. The mixture which shows positive deviation from Raoult's law is : [2020]
1. Chloroethane + Bromoethane 2. Ethanol + Acetone
3. Benzene + Toluene 4. Acetone + Chloroform
16. The freezing point depression constant (K_f) of benzene is $5.12 \text{ K kg mol}^{-1}$. The freezing point depression for the solution of molality 0.078 m containing a non-electrolyte solute in benzene is (rounded off upto two decimal places) [2020]
- 1) 0.60 K 2) 0.20 K 3) 0.80 K 4) 0.40 K
17. The following solutions were prepared by dissolving 10 g of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) in 250 ml of water (P_1), 10 g urea ($\text{CH}_4\text{N}_2\text{O}$) in 250 ml of water (P_2) and 10 g of sucrose ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$) in 250 ml of water (P_3). The right option for the decreasing order of osmotic pressure of these solutions is : [NEET-2021]
1. $P_1 > P_2 > P_3$ 2. $P_2 > P_3 > P_1$ 3. $P_3 > P_1 > P_2$ 4. $P_2 > P_1 > P_3$
18. The correct option for the value of vapour pressure of a solution at 45°C with benzene to octane in molar ratio 3 : 2 is: [NEET-2021]
- [At 45°C vapour pressure of benzene is 280 mm Hg and that of octane is 420 mm Hg. Assume Ideal gas]
- 1) 168 mm of Hg 2) 336 mm of Hg 3) 350 mm of Hg 4) 160 mm of Hg
19. In one molal solution that contains 0.5 mole of a solute, there is [NEET-2022]
- 1) 500 mL of solvent 2) 500 g of solvent
3) 100 mL of solvent 4) 100 g of solvent

NCERT LINE BY LINE QUESTIONS – ANSWERS

(1.)	a	(2.)	c	(3.)	c	(4.)	c	(5.)	b
(6.)	c	(7.)	b	(8.)	a	(9.)	b	(10.)	d
(11.)	a	(12.)	b	(13.)	d	(14.)	b	(15.)	a
(16.)	a	(17.)	c	(18.)	c	(19.)	a	(20.)	c
(21.)	b	(22.)	d	(23.)	c	(24.)	d	(25.)	b
(26.)	d	(27.)	c	(28.)	b	(29.)	c	(30.)	c
(31.)	a	(32.)	c	(33.)	a	(34.)	c	(35.)	b
(36.)	b	(37.)	d	(38.)	d	(39.)	d	(40.)	d
(41.)	b	(42.)	b	(43.)	c	(44.)	b	(45.)	b
(46.)	c	(47.)	a	(48.)	b	(49.)	c	(50.)	c

TOPIC WISE PRACTICE QUESTIONS - ANSWERS

1) 4	2) 4	3) 3	4) 4	5) 2	6) 2	7) 3	8) 4	9) 2	10) 4
11) 4	12) 2	13) 2	14) 2	15) 2	16) 2	17) 2	18) 4	19) 2	20) 4
21) 1	22) 4	23) 2	24) 4	25) 3	26) 4	27) 4	28) 4	29) 2	30) 4
31) 2	32) 1	33) 3	34) 1	35) 3	36) 4	37) 4	38) 3	39) 3	40) 4
41) 1	42) 4	43) 2	44) 4	45) 2	46) 4	47) 4	48) 4	49) 1	50) 1
51) 1	52) 1	53) 4	54) 3	55) 2	56) 3	57) 4	58) 2	59) 3	60) 4
61) 2	62) 2	63) 2	64) 1	65) 3	66) 4	67) 3	68) 4	69) 1	70) 4

NEET PREVIOUS YEARS QUESTIONS-ANSWERS

1) 3	2) 1	3) 1	4) 3	5) 4	6) 1	7) 3	8) 3	9) 1	10) 1
11) 3	12) 3	13) 1	14) 3	15) 2	16) 4	17) 4	18) 2	19) 2	

NCERT LINE BY LINE QUESTIONS – SOLUTIONS

- (1.) (a)
- (2.) (c) Pressure.
Solubility of a solid in liquid does not depend upon pressure since solids and liquids are almost incompressible.
- (3.) (c) Solute – *solvent* Interactions > Solute – *solute* or solvent-solvent interactions
Interactions are high.
So, bonds cannot easily break so that's why vapour decreases so, V.P. decrease negative deviation.
- (5.) (b)
- (6.) (c) $X_{\text{H}_2\text{O}} = 0.02, X_{\text{gas}} = 0.98, P_T = 1.2 \text{ atm}$
Partial Pressure of dry air = $P_T \times \text{M.P of dry air} = 1.2 \times 0.98 = 1.176 \text{ atm}$
- (7.) (b) $A \rightarrow Q, B \rightarrow R, C \rightarrow P$
- (8.) (a) According to Henry's law at constant temperature, the solubility of a gas in a liquid is directly proportional to the pressure of the gas.
- (9.) (b)
- $$\Delta T_f = i \times K_f \times \frac{W_B \times 1000}{M_B \times w_A}$$
- $$\Delta T_f = 3.82, K_f = 1.86$$

$$w_B = 5, M_B = 142, w_A = 45$$

$$i = \frac{\Delta T \times M_B \times w_A}{K_f \times W_B \times 1000} = \frac{3.82 \times 142 \times 45}{1.86 \times 5 \times 1000} = 2.63$$

(11.) (a) According to Boyle Van't Hoff factor $\pi \propto C$ (at constant temperature)

(12.) (b)

$$\alpha = \frac{i-1}{n-1} = \frac{2.74-1}{3-1} = \frac{1.74}{2} = 0.87$$

(14.) (b)

(15.) (a) By Raoult's law

$$P_T = P_P^0 X_P + P_Q^0 X_Q$$

where, $P_P^0 = 80$ torr

$$P_Q^0 = 60 \text{ torr}$$

$$X_P = \frac{3}{5}, X_Q = \frac{2}{5}$$

$$P_T = 80 \times \frac{3}{5} + 60 \times \frac{2}{5} = 72 \text{ torr}$$

(16.) (a) $K_b = \frac{\Delta T_b}{m} = \frac{K}{\text{mol kg}^{-1}}$ or K (molality)

(17.) (c) $\Delta T_f = K_f \times i \times m$

$\Delta T_f \propto i$ (Van't Hoff factor)

Salt	i
KCl	2
$C_6H_{12}O_6$	1
$Al_2(SO_4)_3$	5
K_2SO_4	3

i.e., i is maximum for 5 for $Al_2(SO_4)_3$.

(19.) (a) Minimum boiling azeotrope is formed by solution showing positive deviation, e.g. ethanol + acetone.

(20.) (c) Since the molecular mass of CH_3OH is 32, so, quantity of CH_3OH to prepare 150 ml solution of 2M CH_3OH

$$= \frac{2}{1000} \times 150 \times 32 = 9.6 \text{ g}$$

(21.) (b) According to Raoult's law, the partial pressure of a volatile component of a solution is directly proportional to its mole fraction in solution at any temperature.

(22.) (d) $M_1 V_1 = M_2 V_2$

$$M_1 = 0.02 \text{ M}, V_1 = 4 \text{ L}$$

$$M_2 = ?, V_2 = 5 \text{ L}$$

$$0.02 \times 4 \text{ L} = M_2 \times 5 \text{ L}$$

$$M_2 = 0.016 \text{ M}$$

(25.) (b) Water will move from side (B) to side (A) if pressure greater than osmotic pressure is applied on piston (B) due to reverse osmosis.

- (26.) (d) $P = P_A X_A + P_B X_B$
 $= P_A X_A + P_B (1 - X_A)$
 $= P_A X_A + P_B - P_B X_A$
 $= P_B + X_A (P_A - P_B)$
- (27.) (c) 3.2g
 1.5×10^3 m Aqueous solution of naltorphene, means that 1.5×10^3 mole of Naltorphene is dissolved in 1 kg of water.
Molar mass of Naltorphene = 311 gmol^{-1}
 1.5×10^3 mole of Naltorphene = $1.5 \times 10^3 \times 311 \text{ g} = 0.467 \text{ g}$
Mass of solution = $0.467 + 1000 = 1000.467 \text{ g}$ or, 0.467 g of naltorphene, mass of solution required = 1000.467 g
For 1.5 mg ($1.5 \times 10^3 \text{ g}$) of naltorphene, mass of solution required = $\frac{1000.467}{0.467} \times 1.5 \times 10^{-3} = 3.21 \text{ g}$
- (28.) (b)
- (29.) (c) The value of molal depression constant K_f is constant for a particular solvent, thus, it will be unchanged when molality of the dilute solution is doubled.
- (31.) (a) Depression in freezing point \propto no. of particles $\text{Al}_2(\text{SO}_4)_3$ provides five ions on the ionisation
 $\text{Al}_2(\text{SO}_4)_3 \rightarrow 2\text{Al}^{3+} + 3\text{SO}_4^{2-}$
So, $\text{Al}_2(\text{SO}_4)_3$ have maximum value of depression in freezing point or lowest freezing point.
- (32.) (c) The properties which depend only upon the number of solute particles present in the solution irrespective of their nature are called colligative properties. Osmotic pressure is a colligative property.
- (34.) (d) Entropy of mixing is zero.
- (35.) (b) $\pi = 2.57 \times 10^{-3} \text{ bar}$
 $V = 200 \text{ cm}^3 = 0.200 \text{ litre}$
 $T = 300 \text{ K}$
 $R = 0.083 \text{ L bar K}^{-1} \text{ mol}^{-1}$
use $M_2 = \frac{W_2 RT}{\pi V}$
 $M_2 = \frac{1.26 \text{ g} \times 0.083 \text{ L bar K}^{-1} \text{ mol}^{-1} \times 300 \text{ K}}{2.57 \times 10^{-3} \text{ bar} \times 0.200 \text{ L}}$
 $M_2 = 61022 \text{ gmol}^{-1}$
- (36.) (b) Low atmospheric pressure
At high altitude, the atmospheric pressure is decreased and due to low atmospheric pressure the solubility of oxygen in blood and tissues is reduced. Or Body temperature of human beings remains constant.
- (37.) (d) All of the above.
- (38.) (d) ppm (parts per million.)
- (39.) (d) In relative lowering of vapour pressure, elevation in B.P., depression in freezing point get minimum for high molecular masses but osmotic pressure cannot be lowered as much for high molecular masses.
- (40.) (d) Addition of solute to water decrease the freezing point of water (pure solvent).
 \therefore when 1% lead nitrate (solute) is added to water, the freezing point of water will be below 0°C .
- (41.) (b) $\Delta T_b = i \times K_b \times m$
 $0.216 = 1 \left[\frac{2.16 \times 0.15 \times 1000}{\text{MW} \times 15} \right]$

MW = 100g

- (42.) (b) An ideal solution is as follows:
Volume change (ΔV) of mixing should be zero. Heat change (ΔH) on mixing should be zero. Obey Raoult's law at every range of concentration.

(43.) (c) Molarity
$$= \frac{w \times 1000}{M_w \times V_{\text{sol}} (\text{vol})} = 2$$

$$2 = \frac{w}{63} \times \frac{1000}{250}$$

$$W = \frac{63}{2} \text{ g, mass of acid} \times \frac{70}{100} = \frac{63}{2} \text{ Mass of acid} = 45\text{g}$$



(44.) (b) $1 - \alpha \quad \alpha \quad \alpha$

$$\text{Total} = 1 + \lambda$$

$$i = 1 + \alpha = 1 + 0.2 = 1.2$$

$$\Delta T_f = i \times k_f \times m$$

$$= 1.2 \times 1.86 \times 0.5 = 1.116\text{k} \approx 1.2\text{k}$$

- (46.) (c) For ideal solution

$$\Delta H_{\text{mix}} > 0, \Delta H_{\text{mix}} = +ve$$

$$\Delta V_{\text{mix}} > 0, \Delta V_{\text{mix}} = +ve$$

$$\Delta S_{\text{mix}} > 0$$

(47.) (a)
$$\frac{P_{\text{solvent}}^{\circ} - P_{\text{solution}}}{P_{\text{Solvent}}^{\circ}} = X_{\text{solute}}$$

$$P_A^{\circ} - P_A \text{ (difference in V.P. of pure solvent and solution)} \Rightarrow \text{lowering in V.P. (1)}$$

$$\text{Now, } P_A^{\circ} = \text{V.P. of pure solvent (2)}$$

By dividing (1) and (2) we get,

$$\frac{P_A^{\circ} - P_A}{P_A} = \text{which is relative lowering in V.P.}$$

- (49.) (c)

- (50.) (c) Equal to the rate of crystallization.

Rate of forward reaction (dissolution) = rate of backward reaction (crystallization)

TOPIC WISE PRACTICE QUESTIONS – SOLUTIONS

- (4) For very dil. solution the concentration is expressed in ppm.
- (4) According to Henry's law the mass of a gas dissolved per unit volume of solvent is proportional to the pressure of the gas at constant temperature $m = K p$ i.e. as the solubility increases, value of Henry's law constant decreases. Since CO_2 is most soluble in water among the given set of gases.
- (3) From molarity equation
$$M_1 V_1 + M_2 V_2 = M_3 (V_1 + V_2)$$
$$1 \times 2.5 + 0.5 \times 3 = M_3 \times 5.5$$
$$M_3 = \frac{4}{5.5} = 0.73\text{M}$$
- 4) H_3PO_4 is tribasic so $N = 3M = 3 \times 1\text{N} = 3\text{N}$
- 2) Density = 1.17 gm/cc (given) as $d = \frac{\text{Mass}}{\text{Volume}}$
Volume = 1cc \therefore Mass = $d = 1.17\text{g}$

$$\text{Molarity} = \frac{\text{No. of moles}}{\text{Volume in litre}} = \frac{1.17 \times 1000}{36.5 \times 1} = \frac{1170}{36.5} = 32.05\text{M}$$

6. 2) Relation between molality and mole fraction is

$$m = \frac{1000 \times x_2}{x_1 M_1} = \frac{1000 \times 0.2}{0.8 \times 78} = 3.2$$

$$\text{Thus, } X(m) = 3.2$$

7. 3) From molarity equation

$$M_1 V_1 + M_2 V_2 = M_3 (V_1 + V_2)$$

$$1 \times 2.5 + 0.5 \times 3 = M_3 \times 5.5$$

$$M_3 = \frac{4}{5.5} = 0.73\text{M}$$

8. 4) In acidic medium, $\text{K}_2\text{Cr}_2\text{O}_7$ undergo reduction as follows:



Change in oxidation states = $6 - 3 = 3$

\therefore Net change = $2 \times 3 = 6$ [Two Cr atoms are involved]

\therefore Eq wt. per unit of $\text{K}_2\text{Cr}_2\text{O}_7 = M/6$

9. 2) One molal solution means one mole of solute is present in 1 kg (1000 g) solvent i.e., mole of solute = 1

$$\text{Mole of solvent (H}_2\text{O)} = \frac{1000\text{g}}{18\text{g}} = \frac{1000}{18}$$

$$\text{Mole fraction of solute} = \frac{1}{\left(1 + \frac{1000}{18}\right)} = \frac{18}{1008} = 0.018$$

10. 4) $\text{ppm} = \frac{\text{Mass of solute}}{\text{Mass of solution}} \times 10^6$

$$\therefore \text{ppm} = \frac{10}{10^6} \times 10^6 = 10 \text{ ppm}$$

11. 4) All get dissolved with the evolution of heat.

12. 2) $\text{Molarity} = \frac{\text{No. of moles of solute}}{\text{Volume in litres}} = \frac{5}{2.5} = 2\text{M}$

13. 2) An increase in temperature of the solution increases the solubility of a solid solute. The amount of solute that dissolve depends on what type of solute it is. For solids and liquid solutes, changes in pressure have practically no effect on solubility.

14. 2) $P_{\text{solution}} = P_{\text{solution}}^0 x_{\text{solvent}}$; $\frac{P^0 - P}{P^0} = x_{\text{solute}}$

15. 2) $\frac{P^0 - P_s}{P^0} = \text{Mole fraction of solute} = \lambda_2$

16. 2)

Given $P_1 = 19 \text{ mm Hg}$, $P_2 = 760 \text{ mm Hg}$;

$\Delta H_{\text{vap.}} = 40670 \text{ J/mol}$

Applying Clausius-Clapeyron's equation

$$\log \frac{P_2}{P_1} = \frac{\Delta H_{\text{vap}}}{2.303 \times R} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$$

$$\text{or } \log \frac{760}{19} = \frac{40670}{2.303 \times 8.3} \left(\frac{373 - T_1}{T_1 \times 373} \right)$$

on solving, we get $T_1 = 291.4 \text{ K}$

17. 2) $P = P_A^0 x_A + P_B^0 x_B = P_B^0 + x_A (P_A^0 - P_B^0) [\because x_B = 1 - x_A]$

18. 4) $\frac{P^0 - P}{P^0} = \frac{10}{50} = \frac{n}{n+12} \therefore n = 3$

19. 2) Equimolar solutions of normal solutes in the same solvent will have the same b. p and same f. p.

20. 4) Solutions in cases I and II are saturated and that in III is unsaturated.

21. 1) Given V. $P_P = 80 \text{ torr}$

V. $P_P = 80 \text{ torr}$

$$P_{\text{total}} = V.P_P \times x_P + V.P_Q \times x_Q = \left[80 \times \frac{3}{5} + 60 \times \frac{2}{5} \right] = 16 \times 3 + 12 \times 2$$

$$P_{\text{total}} = 48 + 24 = 72 \text{ torr}$$

22. 4) A solution containing A and B components shows negative deviation when A-A and B-B interactions are weaker than that of A-B interactions. For such solutions.

$\Delta H = -\text{ve}$ and $\Delta V = -\text{ve}$

23. 2) Number of solute particles going into solution will be equal to the solute particles separating out and a state of dynamic equilibrium is reached.

solute + solvent \rightleftharpoons solution

i.e., rate of dissolution = rate of crystallization

24. 4) $\frac{P^0 - P}{P^0} = \frac{n_2}{n_1 + n_2}$

$$\frac{640 - 600}{640} = \frac{25.5 / m}{39 / 78}$$

$$m = \frac{640 \times 78 \times 2.5}{39 \times 40} = 80$$

25. 3) For ideal solution, $\Delta V_{\text{mixing}} = 0$ and $\Delta H_{\text{mixing}} = 0$


26. 4) Gaseous densities of ethanol and dimethyl ether would be same at same temperature and pressure. The heat of vaporisation, V.P. and b.pts will differ due to H-bonding in ethanol.

27. 4)

28. 4) Addition of a solute increases the boiling point of solution.

29. 2) Solvent having high cryoscopic constant (camphor) can be used in determination of molecular masses of organic compounds like naphthalene, anthracene etc., by cryoscopic method.

30. 4) Boiling temperature is a temperature at which vapour pressure is equal to the atmospheric pressure when external pressure is equal to 1 atm.



The number of ions decreases in the ratio of 4 to 3, and so also the colligative property.

32. 1) Lowering is always positive

33. 3) $\frac{P^\circ - P_s}{P^\circ} = \frac{n}{N} = \frac{w}{m} \times \frac{M}{W}$; $0.1 = \frac{12}{m} \times \frac{18}{108}$; $m = \frac{12 \times 18}{0.1 \times 108} = 20$
34. 1) Osmosis occurs from dilute solution to concentrated solution, i.e., exosmosis.
35. 3) Osmotic pressure \propto no. of ions
(NH_4)₃ PO₄ gives maximum ions. Hence its osmotic pressure is maximum
36. 4) $\pi = CRT \Rightarrow 2.5 = C \times 0.0821 \times 297$
 $\therefore C = 0.1025 \text{ mol L}^{-1}$
37. 4) $\pi = MRT$
i) $\pi = RT$; ii) $\pi = RT$; iii) $\pi = RT$ (mix has concentration = 1M)
38. 3) Osmotic pressure is a colligative property
39. 3) B.P. \propto moles of non-volatile solute.
40. 4) Liquid solution \square solid solvent
41. 1) $\Delta T_f = i \times K_f \times m$
Van't Hoff factor, $i = 2$ for NaCl, $m = 0.01$ hence $\Delta T_f = 0.02K_f$ which is maximum in the present case. Hence ΔT_f is maximum or freezing point is minimum.
42. 4) $0.465 = 1.86 \times \left[\frac{1.8/x}{40} \times 10^3 \right]$
 $x = 180 \text{ g/mol} \therefore \text{C}_6\text{H}_{12}\text{O}_6$
43. 2) $T_f = \frac{\Delta H}{\Delta S}$
44. 4) Depression in freezing point, $\Delta T_f = iK_f m$. The value of van't Hoff factor (i) is minimum for the glucose, which is a non-electrolyte. Hence, aqueous solution of glucose has highest freezing point.
45. 2) $\text{KCl} \rightleftharpoons \text{K}^+ + \text{Cl}^-$
 $1-\alpha \quad \alpha \quad \alpha$
Total number of particles at e.g.
 $1-\alpha + \alpha + \alpha = 1 + \alpha = 1 + 0.5$ $\Delta T_f = i k_f m$
 $\Delta T_f = (1 + 0.5) \times 1.86 \times 0.5 = 1.395$
 $\therefore T_f = 273 - 1.395 = 271.60$
46. 4) $i = 1 + \alpha \Rightarrow 1.25$
 $\Delta T_f = K_f . m . i \Rightarrow 1.86 \times 0.2 \times 1.25$ or $\Delta T_f = 0.465$ or $T_f = -0.465^\circ \text{C}$
47. 4) Depression in freezing point = $K_f m$ (as $K_f \uparrow, \Delta T_f \uparrow$)
 K_f depends upon the solvent
48. 4)
 $t=0 \quad \text{AB}_3 \rightleftharpoons \text{A}^{3+} + 3\text{B}^-$
 $\quad \quad \quad 1 \quad \quad \quad 0 \quad \quad 0$
 $t=t_{\text{eq}} \quad 1-\alpha \quad \quad \alpha \quad \quad \alpha$
Total $1 + \alpha + \alpha + 3\alpha = 1 + 3\alpha$
 $\therefore T - 100 = 0.1 (1 + 3 \times 0.9) \times 0.52$
 $T = 373.19 \text{ K}$
49. 1) In osmosis, solvent molecules move from lower concentration (higher vapour pressure) to higher concentration (Lower vapour pressure)
50. 1) $\pi \propto i$ [Equal concentration]
Aluminium sulphate $\text{Al}_2(\text{SO}_4)_3$ ($i = 5$),
Barium chloride BaCl_2 ($i = 3$),

51. (1) Concentration of particles in CaCl_2 solution will be maximum as $i = 3$ for CaCl_2 and $i = 2$ for KCl .

Glucose and Urea do not dissociate into ions, as they are nonelectrolytes.

52. 1) $7.4 = n \times 0.0821 \times 300$
 $\therefore n = 0.3$

53. 4) Addition of solute to water decreases the freezing point of water (pure solvent).
 \therefore When 1% lead nitrate (solute) is added to water, the freezing point of water will be below 0°C .

54. 3) $\Delta T_b = \frac{K_b \times 100 \times 100}{10 \times 1000} = \Delta T_b$

55. 2) $M = \frac{K_b \times w \times 1000}{\Delta T_b \times W} = \frac{2.16 \times 0.11 \times 1000}{0.1 \times 15} = 158.4$

56. 3) $K_b = \frac{0.1 \times 180 \times 100}{1.8 \times 1000} = 1\text{K/m}$

57. 4) $0.186 = 1.86 \times m; \therefore m = 0.1;$
 $\therefore \Delta T_b = 0.512 \times 0.1 = 0.0512^\circ\text{C}$

58. 2) $\pi V = CRT$

$$\frac{\pi_1}{\pi_2} = \frac{C_1 T_1}{C_2 T_2}$$

$$\pi_1 = P, \pi_2 = 2\text{atm}, C_1 = C, C_2 = \frac{C}{2}$$

$$T_1 = 600\text{K}, T_2 = 700\text{K}$$

$$\frac{P}{2} = \frac{2 \times C \times 600}{C \times 700}; P = \frac{24}{7}$$

59. 3)

60. 4) Osmosis is a spontaneous process ($\Delta G < 0$) which takes place with ($\Delta S < 0$) (due to transfer of solvent in 70 solution)

61. 2) $\Delta T_b = K_b \times i \times m$

Where ΔT_b = Elevation in boiling point

K_b = molal elevation constant


i = vant Hoff factor

$\therefore \Delta T_b \propto \text{molality}$


62. 2) Benzoic acid exists as dimer in benzene.

63. 2) Blood cells neither swell nor shrink in isotonic solution. As isotonic solutions have equal concentration therefore there is no flow of solvent occurs and hence solvent neither enters nor flow out of the blood cells.

64. 1) The solution which provide same number of ions are isotonic.



Total ions produced = 3




Total ions produced = 3

$\therefore 0.1\text{M } \text{Ca}(\text{NO}_3)_2$ and $0.1\text{M } \text{Na}_2\text{SO}_4$ are isotonic

65. 3) If compound dissociates in solvent $i > 1$ and on association $i < 1$.

66. 4)

67. 3)



Mol. before dissociation 1 0 0

Mol. after dissociation 1 - α 2α 1α



$$i = 1 - \alpha + 2\alpha + \alpha = 1 + 2\alpha$$

68. 4) Osmotic pressure is directly proportional to the number of particles in the solution. KCl will dissociate completely in the solution, while CH_3COOH and sucrose are weak electrolyte.

Hence correct order will be :



69. 1)
 $\text{K}_4[\text{Fe}(\text{CN})_6]$ and $\text{Al}_2(\text{SO}_4)_3$ both dissociates to give 5 ions or $i = 5$



70. 4) All the statements are correct

NEET PREVIOUS YEARS QUESTIONS-EXPLANATIONS

1. 3) K_f (molal depression constant) only depends on the nature of the solvent and is independent of the concentration of the solution.

2. 1) Let us consider that A is benzene and B is toluene 1 : 1 molar mixture of A and B

$$\therefore x_A = \frac{1}{2} \text{ and } x_B = \frac{1}{2}$$

$$\text{Total pressure of solution (P)} = P_A^0 x_A + P_B^0 x_B$$

$$P = 12.8 \times \frac{1}{2} + 3.85 \times \frac{1}{2} = 8.325 \text{ kPa} ; \quad Y_A = \frac{P_A^0 x_A}{P} = \frac{12.8 \times \frac{1}{2}}{8.325} = 0.768$$

$$\therefore y_B = 1 - y_A = 1 - 0.768 = 0.232$$

so, the vapour will contain higher percentage of benzene.

3. 1)

$$\left(\frac{P^0 - P_s}{P^0} \right) = \frac{n}{N} = \frac{W_1}{M_1} \times \frac{M_2}{W_2}$$

Where, W_1 = wt of solute

W_2 = wt of solvent

M_1 = Mass of solute

M_2 = Mass of solvent

at 100°C , $P^0 = 760 \text{ mm}$

$$\frac{760 - 732}{760} = \frac{6.5 \times 18}{M_1 \times 100}$$

$$M_1 = 31.75 \text{ g mol}^{-1}$$


$$\Delta T_b = m \times K_b = \frac{W_1 \times 1000}{M_1 \times W_2} \times K_b$$

$$\Delta T_b = \frac{0.52 \times 6.5 \times 1000}{31.75 \times 100} = 1.06^\circ\text{C}$$

boiling point of solution

$$= 100^\circ\text{C} + 1.06^\circ\text{C} = 101^\circ\text{C}$$

4. 3) $\text{K}_4[\text{Fe}(\text{CN})_6] \rightleftharpoons 4\text{K}^+ + [\text{Fe}(\text{CN})_6]^-$



\therefore van't Hoff factor is 5 for both



5. 4) $\Delta T_b = iK_b m$

Given $(\Delta T_b)_x > (\Delta T_b)_y$

$\therefore i_x K_b m > i_y K_b m$ (K_b is same for same solvent)

$i_x > i_y$ So, x is undergoing dissociation in water

6. 1) For an ideal solution $\Delta S_{\text{mix}} > 0$

7. 3) Colligative properties \propto no. of particles since $\text{Al}_2(\text{SO}_4)_3$ contains maximum number of particles,

hence will have the largest value of freezing point depression.

8. 3) For an ideal solution, $\Delta H_{\text{mix}} = 0$

9. 1) Maximum boiling azeotrope are formed by solutions which show negative deviation from ideal behaviour. Water + Nitric acid shows negative deviation.

10. 1)

If, intermolecular attractive forces between **A - A**

and **B - B** are **stronger** than those **A - B** then

it show positive deviation from ideal behaviour.

11. 3)

$X_{\text{H}_2\text{O}} = 0.02$

$\therefore X_{\text{gas}} = 0.98$

$P_{\text{total}} = 1.2 \text{ atm}$

partial pressure of dry-air = $P_T \times$ mole fraction of dry-air

Partial pressure of dry-air = $1.2 \text{ atm} \times 0.98$

= 1.176 atm.

12. 3)

13. 1) Assuming dilute solution,

$$\frac{P_0 - P_s}{P_s} \approx \frac{P_0 - P_s}{P_0} = \frac{n_{\text{solute}}}{n_{\text{solvent}}}$$

Let $P_0 = 100$, V.P reduced to 80%, $\therefore P_s = 80$

$$\frac{100 - 80}{100} = \frac{8 / m}{114 / 114}$$

$m = 40$

14. 3) Isotonic solutions have same osmotic pressure.

15. 2) Ethanol + Acetone mixture shows positive deviation from Raoult's law

16. 4) $\Delta'_F = K_f \times m = 5.12 \times 0.078$; Depression in = 0.4K ; Freezing point

17. 4) $\pi = i.C.R.T$

$\pi \propto \frac{1}{\text{molecular weight}} ; \quad P_2 > P_1 > P_3$

18. 2) $\frac{C_6H_6}{n_1 = 3}$ $\frac{C_8H_{18}}{n_2 = 2}$

$P_1 = 280 \text{ mm}$

$P_2 = 420 \text{ mm}$

$$P = P_1X_1 + P_2X_2 = 280 \times \frac{3}{5} + 420 \times \frac{2}{5} = 168 + 168 = 336 \text{ mm of Hg}$$

19. $\text{Molality} = n \times \frac{1000}{\text{weight of solvent}}$

$$1 = 0.5 \times \frac{1000}{w} \Rightarrow w = 500 \text{ g}$$

**CLICK HERE TO
DOWNLOAD
LINE BY LINE QUESTIONS
CLASS 12 ALL SUBJECTS**





JOIN OUR WHATSAPP GROUPS

FOR FREE EDUCATIONAL
RESOURCES





JOIN SCHOOL OF EDUCATORS WHATSAPP GROUPS FOR FREE EDUCATIONAL RESOURCES

We are thrilled to introduce the School of Educators WhatsApp Group, a platform designed exclusively for educators to enhance your teaching & Learning experience and learning outcomes. Here are some of the key benefits you can expect from joining our group:

BENEFITS OF SOE WHATSAPP GROUPS

- **Abundance of Content:** Members gain access to an extensive repository of educational materials tailored to their class level. This includes various formats such as PDFs, Word files, PowerPoint presentations, lesson plans, worksheets, practical tips, viva questions, reference books, smart content, curriculum details, syllabus, marking schemes, exam patterns, and blueprints. This rich assortment of resources enhances teaching and learning experiences.
- **Immediate Doubt Resolution:** The group facilitates quick clarification of doubts. Members can seek assistance by sending messages, and experts promptly respond to queries. This real-time interaction fosters a supportive learning environment where educators and students can exchange knowledge and address concerns effectively.
- **Access to Previous Years' Question Papers and Topper Answers:** The group provides access to previous years' question papers (PYQ) and exemplary answer scripts of toppers. This resource is invaluable for exam preparation, allowing individuals to familiarize themselves with the exam format, gain insights into scoring techniques, and enhance their performance in assessments.

- **Free and Unlimited Resources:** Members enjoy the benefit of accessing an array of educational resources without any cost restrictions. Whether its study materials, teaching aids, or assessment tools, the group offers an abundance of resources tailored to individual needs. This accessibility ensures that educators and students have ample support in their academic endeavors without financial constraints.
- **Instant Access to Educational Content:** SOE WhatsApp groups are a platform where teachers can access a wide range of educational content instantly. This includes study materials, notes, sample papers, reference materials, and relevant links shared by group members and moderators.
- **Timely Updates and Reminders:** SOE WhatsApp groups serve as a source of timely updates and reminders about important dates, exam schedules, syllabus changes, and academic events. Teachers can stay informed and well-prepared for upcoming assessments and activities.
- **Interactive Learning Environment:** Teachers can engage in discussions, ask questions, and seek clarifications within the group, creating an interactive learning environment. This fosters collaboration, peer learning, and knowledge sharing among group members, enhancing understanding and retention of concepts.
- **Access to Expert Guidance:** SOE WhatsApp groups are moderated by subject matter experts, teachers, or experienced educators can benefit from their guidance, expertise, and insights on various academic topics, exam strategies, and study techniques.

Join the School of Educators WhatsApp Group today and unlock a world of resources, support, and collaboration to take your teaching to new heights. To join, simply click on the group links provided below or send a message to +91-95208-77777 expressing your interest.

**Together, let's empower ourselves & Our Students and
inspire the next generation of learners.**

**Best Regards,
Team
School of Educators**

Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

Kindergarten to Class XII (For Teachers Only)



[Click Here to Join](#)

Class 1



[Click Here to Join](#)

Class 2



[Click Here to Join](#)

Class 3



[Click Here to Join](#)

Class 4



[Click Here to Join](#)

Class 5



[Click Here to Join](#)

Class 6



[Click Here to Join](#)

Class 7



[Click Here to Join](#)

Class 8



[Click Here to Join](#)

Class 9



[Click Here to Join](#)

Class 10



[Click Here to Join](#)

Class 11 (Science)



[Click Here to Join](#)

Class 11 (Humanities)



[Click Here to Join](#)

Class 11 (Commerce)



[Click Here to Join](#)

Class 12 (Science)



[Click Here to Join](#)

Class 12 (Humanities)



[Click Here to Join](#)

Class 12 (Commerce)



[Click Here to Join](#)

Kindergarten

Subject Wise Secondary and Senior Secondary Groups (IX & X For Teachers Only) Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi-A



[Click Here to Join](#)

IT Code-402



[Click Here to Join](#)

Hindi-B



[Click Here to Join](#)

Artificial Intelligence

Senior Secondary Groups (XI & XII For Teachers Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History



[Click Here to Join](#)

Geography



[Click Here to Join](#)

Sociology



[Click Here to Join](#)

Hindi Elective



[Click Here to Join](#)

Hindi Core



[Click Here to Join](#)

Home Science



[Click Here to Join](#)

Sanskrit



[Click Here to Join](#)

Psychology



[Click Here to Join](#)

Political Science



[Click Here to Join](#)

Painting



[Click Here to Join](#)

Vocal Music



[Click Here to Join](#)

Comp. Science



[Click Here to Join](#)

IP



[Click Here to Join](#)

Physical Education



[Click Here to Join](#)

APP. Mathematics



[Click Here to Join](#)

Legal Studies



[Click Here to Join](#)

Entrepreneurship



[Click Here to Join](#)

French



[Click Here to Join](#)

IT



[Click Here to Join](#)

Artificial Intelligence

Other Important Groups (For Teachers & Principal's)



[Click Here to Join](#)

Principal's Group



[Click Here to Join](#)

Teachers Jobs



[Click Here to Join](#)

IIT/NEET

Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

Kindergarten to Class XII (For Students Only)



[Click Here to Join](#)

Class 1



[Click Here to Join](#)

Class 2



[Click Here to Join](#)

Class 3



[Click Here to Join](#)

Class 4



[Click Here to Join](#)

Class 5



[Click Here to Join](#)

Class 6



[Click Here to Join](#)

Class 7



[Click Here to Join](#)

Class 8



[Click Here to Join](#)

Class 9



[Click Here to Join](#)

Class 10



[Click Here to Join](#)

Class 11 (Science)



[Click Here to Join](#)

Class 11 (Humanities)



[Click Here to Join](#)

Class 11 (Commerce)



[Click Here to Join](#)

Class 12 (Science)



[Click Here to Join](#)

Class 12 (Humanities)



[Click Here to Join](#)

Class 12 (Commerce)



[Click Here to Join](#)

**Artificial Intelligence
(VI TO VIII)**

Subject Wise Secondary and Senior Secondary Groups (IX & X For Students Only) Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi



[Click Here to Join](#)

IT Code



[Click Here to Join](#)

Artificial Intelligence

Senior Secondary Groups (XI & XII For Students Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History



[Click Here to Join](#)

Geography



[Click Here to Join](#)

Sociology



[Click Here to Join](#)

Hindi Elective



[Click Here to Join](#)

Hindi Core



[Click Here to Join](#)

Home Science



[Click Here to Join](#)

Sanskrit



[Click Here to Join](#)

Psychology



[Click Here to Join](#)

Political Science



[Click Here to Join](#)

Painting



[Click Here to Join](#)

Music



[Click Here to Join](#)

Comp. Science



[Click Here to Join](#)

IP



[Click Here to Join](#)

Physical Education



[Click Here to Join](#)

APP. Mathematics



[Click Here to Join](#)

Legal Studies



[Click Here to Join](#)

Entrepreneurship



[Click Here to Join](#)

French



[Click Here to Join](#)

IT



[Click Here to Join](#)

AI



[Click Here to Join](#)

IIT/NEET



[Click Here to Join](#)

CUET

Groups Rules & Regulations:

To maximize the benefits of these WhatsApp groups, follow these guidelines:

1. Share your valuable resources with the group.
2. Help your fellow educators by answering their queries.
3. Watch and engage with shared videos in the group.
4. Distribute WhatsApp group resources among your students.
5. Encourage your colleagues to join these groups.

Additional notes:

1. Avoid posting messages between 9 PM and 7 AM.
2. After sharing resources with students, consider deleting outdated data if necessary.
3. It's a NO Nuisance groups, single nuisance and you will be removed.
 - No introductions.
 - No greetings or wish messages.
 - No personal chats or messages.
 - No spam. Or voice calls
 - Share and seek learning resources only.

Please only share and request learning resources. For assistance, contact the helpline via WhatsApp: +91-95208-77777.

Join Premium WhatsApp Groups Ultimate Educational Resources!!

Join our premium groups and just Rs. 1000 and gain access to all our exclusive materials for the entire academic year. Whether you're a student in Class IX, X, XI, or XII, or a teacher for these grades, Artham Resources provides the ultimate tools to enhance learning. Pay now to delve into a world of premium educational content!

[Click here for more details](#)



Click Here to Join

Class 9



Click Here to Join

Class 10



Click Here to Join

Class 11



Click Here to Join

Class 12

📢 Don't Miss Out! Elevate your academic journey with top-notch study materials and secure your path to top scores! Revolutionize your study routine and reach your academic goals with our comprehensive resources. Join now and set yourself up for success! 🇧🇩🌟

Best Wishes,

Team

School of Educators & Artham Resources

SKILL MODULES BEING OFFERED IN MIDDLE SCHOOL



Artificial Intelligence



Beauty & Wellness




Design Thinking & Innovation



Financial Literacy



Handicrafts



Information Technology



Marketing/Commercial Application




Mass Media - Being Media Literate




Travel & Tourism



Coding




Data Science (Class VIII only)




Augmented Reality / Virtual Reality



Digital Citizenship




Life Cycle of Medicine & Vaccine



Things you should know about keeping Medicines at home




What to do when Doctor is not around



Humanity & Covid-19



Blue Pottery



Pottery



Block Printing




Food



Food Preservation




Baking



Herbal Heritage



Khadi




Mask Making




Mass Media



Making of a Graphic Novel



Kashmiri Embroidery




Embroidery




Rockets



Satellites




Application of Satellites



Photography

SKILL SUBJECTS AT SECONDARY LEVEL (CLASSES IX – X)



Retail



Information Technology




Security




Automotive




Introduction To Financial Markets



Introduction To Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking & Insurance



Marketing & Sales



Health Care



Apparel



Multi Media



Multi Skill Foundation Course



Artificial Intelligence



Physical Activity Trainer



Data Science



Electronics & Hardware (NEW)



Foundation Skills For Sciences (Pharmaceutical & Biotechnology)(NEW)



Design Thinking & Innovation (NEW)

SKILL SUBJECTS AT SR. SEC. LEVEL (CLASSES XI – XII)



Retail



Information Technology



Web Application



Automotive



Financial Markets Management



Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking




Marketing



Health Care



Insurance




Horticulture



Typography & Comp.
Application



Geospatial Technology



Electrical Technology



Electronic Technology



Multi-Media




Taxation



Cost Accounting



Office Procedures & Practices



Shorthand (English)




Shorthand (Hindi)



Air-Conditioning & Refrigeration



Medical Diagnostics



Textile Design




Design



Salesmanship



Business Administration



Food Nutrition & Dietetics



Mass Media Studies



Library & Information Science



Fashion Studies



Applied Mathematics



Yoga



Early Childhood Care & Education




Artificial Intelligence



Data Science



Physical Activity Trainer(new)



Land Transportation Associate (NEW)



Electronics & Hardware (NEW)



Design Thinking & Innovation (NEW)

